

11/15/00
JC950 U.S. PTO

11/16/00 not for
IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

A

Applicant: Uckun
Docket: 12152.109US01
Title: VANADIUM COMPOUNDS AS ANTI-ANGIOGENIC AGENTS

JC948 U.S. PTO
09/713780
11/15/00

CERTIFICATE UNDER 37 CFR 1.10

'Express Mail' mailing label number: EL649973516US

Date of Deposit: November 15, 2000

I hereby certify that this paper or fee is being deposited with the United States Postal Service 'Express Mail Post Office To Addressee' service under 37 CFR 1.10 and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231.

By:

Name: Denise M. Kettelberger

BOX PATENT APPLICATION

Assistant Commissioner for Patents

Washington, D.C. 20231

Sir:

We are transmitting herewith the attached:

- ☒ Transmittal sheet, in duplicate, containing Certificate under 37 CFR 1.10.
- ☒ Utility Patent Application: Spec. 24 pgs; 24 claims; Abstract 1 pgs.:
- ☒ Small entity status will be established at a later date
- ☒ An unsigned Combined Declaration and Power of Attorney
- ☒ Other: Bibliographic Sheet
- ☒ Return postcard
- ☒ **PAYMENT OF THE FILING FEE IS BEING DEFERRED.**

MERCHANT & GOULD P.C.

P.O. Box 2903, Minneapolis, MN 55402-0903

(612) 332-5300

By:

Name: Denise M. Kettelberger

Reg. No.: 33,924

Initials: DMK:mcp



23552

PATENT TRADEMARK OFFICE

Inventor Information

Inventor One Given Name :: Fatih M.
Family Name :: Uckun
Name Suffix ::
Postal Address Line One :: 12590 Ethan Avenue North
Postal Address Line Two ::
City :: White Bear Lake
State/Province :: Minnesota
Country :: USA
Postal or Zip Code :: 55110
City of Residence :: White Bear Lake
State/Prov. of Residence :: Minnesota
Country of Residence :: USA
Citizenship :: USA

Given name of Applicant ::
Family Name ::
Name Suffix ::
Authority under 1.42 ::
Authority under 1.43 ::
Authority under 1.47 ::
Postal Address Line One ::
Postal Address Line Two ::
City ::
State/Province ::
Country ::
Postal or Zip Code ::
City of Residence ::
State/Prov. of Residence ::
Country of Residence ::
Citizenship ::

Correspondence Information

Correspondence Customer Number :: 23552
Name Line One :: Denise M. Kettelberger, Ph.D.
Name Line Two ::
Address Line One :: Merchant & Gould P.C.
Address Line Two :: P.O. Box 2903
City :: Minneapolis
State/Province :: Minnesota
Country :: USA
Postal or Zip Code :: 55402-0903
Telephone :: 612/371-5268
Fax :: 612/332-9081
E-Mail :: dkettelberger@merchant-gould.com

Application Information

Title Line One :: VANADIUM COMPOUNDS AS ANTI-ANGIOGENIC AGENTS

Total Drawing Sheets ::

Formal Drawings ::

Application Type :: Patent

Docket Number :: 12152.109US01

Representative Information

Representative Customer Number :: 23552

Registration Number One :: 33,924



00577-0923260

VANADIUM COMPOUNDS AS ANTI-ANGIOGENIC AGENTS

Background of the Invention

Tissue growth is intimately associated with blood supply. Non-vascularized tissue is limited in size, often being smaller than one to two millimeters in diameter or thickness. Therefore, inhibiting blood supply to tissue represents one target point for limiting tissue growth and possibly tissue viability.

The ability to inhibit blood supply has been shown to play a pivotal role in the progression, invasive and metastatic growth of malignant tumors (Folman, *Nat Med*, 1:27, 1995; Folkman et al, *Science*, 235:442, 1985; Gimbrane et al, *J Exp Med*, 136:261, 1972). Further, inhibition of angiogenesis new vessel formation), has been shown to result in tumor dormancy or regression and to prevent metastasis (Folkman, *Ann Surg*, 175:409, 1972; Taylor et al, *Nature*, 297:307, 1982).

Another possible target for inhibiting tissue growth is by inhibiting cell proliferation. All proliferating eucaryotic cells must undergo mitosis before separating into two new cells. Mitosis is a process in which the parent or replicating cell undergoes a series of molecular events that results in the formation of two nuclei in the place of one.

The polar mitotic spindle is critical to the separation of the replicated chromosomes and formation of the two nuclei in the mitotic process. For mitosis to proceed normally, cells must properly form a bipolar mitotic spindle with bivalent chromosomes properly attached to each pole of the spindle (Gorbsky et al, *Bioessays*, 19: 193-197, 1997; Hardwick, K. G., *Trends Genet.*, 14: 1-4, 1998). Cells which do not form a correct mitotic spindle arrest indefinitely in the metaphase stage of mitosis or progress into apoptosis. Several proteins identified in yeast and mammals have been implicated in this process, including MAD1 (mitotic arrest deficient), MAD2, and MAD3 (Li et al, *Cell*, 66: 519-531, 1991 (*published erratum appears in Cell*, 79(2), following p388)), BUB1 (budding uninhibited by benzimidazole), BUB2 and BUB3 (Hoyt et al, *Cell*, 66: 507-517, 1991). Mammalian counterparts for these proteins include HsMAD2 (Li et al, *Supra* and hBUB1 (Cahill et al, *Nature*, 392: 300-303, 1998).

Revascularization of obstructed coronary arteries by percutaneous transluminal coronary angioplasty (PTCA) has become an integral component of front-line treatment programs for patients with ischemic heart disease (Vaitkus, P.T., 1995, *Coron. Artery Dis.*, 6:429-439). Although acute complications of PTCA have markedly declined with
5 optimized use of anticoagulants, antispasmodic agents, and intravascular stents, the incidence of coronary artery restenosis has remained at 30%-50% and represents the major obstacle to a more successful outcome of PTCA (Landzberg, et. al., 1997, *Prog. Cardiovascular Diseases*, 39:361-298). Therefore, the development of effective
10 strategies for restenosis prophylaxis has become a focal point for translational cardiovascular research.

The pathogenesis of restenosis has been compared to an exaggerated wound healing response with migration of smooth muscle cells from the media to the intima of the revascularized coronary artery where they proliferate and cause an obstructive neointimal hyperplasia (Ueda et al., 1995, *Coron. Artery Dis.*, 6:71-81).
15 Many factors contribute to the development of restenosis, including vascular injury, platelet aggregation, procedural factors, inflammation, and mitogenic stimulation of migration and proliferation of smooth muscle cells. The relative contribution of any one of these factors remains unclear.

Pharmacological approaches to prevent restenosis include antiplatelet
20 and antithrombotic agents, anti-inflammatory drugs, growth factor antagonists, vasodilators, antiproliferatives, antineoplastics, photochemotherapy, and lipid lowering agents. Some growth factor antagonists have also been studied for effects on restenosis.

Inhibition of vascular smooth muscle cell proliferation by a platelet derived growth factor (PDGF)-antagonist has generated promising results in preclinical
25 as well as early clinical studies, thereby confirming the biologic importance of vascular smooth muscle cells in the pathophysiology of restenosis (Mullins et al., 1994, *Arterioscler. Thromb.*, 14:1047-1055).

Considerable efforts are underway to develop new anti-angiogenic and
30 anti-mitotic agents for use as therapies in the treatment of tumor growth and spread.

Accordingly, there is a need for the analysis of novel, effective anti-angiogenic and anti-mitotic agents that target tumor growth.

Vanadocene dichloride (VDC) has been shown to arrest tumor cells growth (Kopf-Maier, et al, *J Cancer Res. Clin. Oncol.*, 106: 44-52, 1983). The
 5 oxovanadium compound, $\text{VO(Phen)(H}_2\text{O)}_2\text{](SO}_4\text{)}$, has been shown to be an active agent against pharyngonasal cancer as determined by a single assay (Sakurai, et. al, *BBRC*, 206: 133, 1995). Vanadium compounds, including vanadocenes, have also been demonstrated to induce apoptosis in cancer cells (Uckun et al., WO 00/35930).

Against this backdrop the present invention has been developed.

10

Summary of the Invention

It has been now found that vanadium compounds, for example vanadocenes and oxovanadium compounds, are poent agents for inhibiting angiogenesis. In particular, the vanadium compounds have been found to exhibit dual
 15 function inhibiting both angiogenesis and mitosis. Pharmaceutical compositions containing these vanadium compounds are thus useful in methods to inhibit angiogenesis and mitosis, for example, in the treatment diabetic retinopathy, hemangiomas, cancers with abnormal blood vessel supply, restenosis following vascular injury, and the like.

20

Detailed Description of the Preferred Embodiment

The present invention relates to vanadium compounds, including, for example, vanadium cyclopentadienyl complexes (vanadocenes), such as vandocene acetylacetonate (VDacac) and oxovanadium compounds, and the finding that these vanadium compounds have potent anti-angiogenic and anti-mitotic activities.

25

Vanadium is a physiologically essential element that can be found in both anionic and cationic forms with oxidation states ranging from -3 to $+5$ (I-V). This versatility provides unique properties to vanadium complexes. In particular, the catonic form of vanadium complexes having an oxidation state of $+4$ (IV) has been shown to function as a modulator of cellular redox potential, regulate enzymatic phosphorylation,

and exert pleiotropic effects in multiple biological systems by catalyzing the generation of reactive oxygen species (ROS). Besides the ability of vanadium metal to assume various oxidation states, its coordination chemistry also plays a key role in its interactions with various biomolecules. In particular, it is demonstrated herein that vanadium compounds, such as vanadium cyclopentadienyl complexes, oxovanadium complexes, or derivatives thereof, exhibit anti-angiogenic and anti-mitotic properties.

Definitions

The following terms and phrases as used herein have the noted definitions, unless otherwise described:

"Halo" is fluoro, chloro, bromo, or iodo.

"Alkyl", "alkoxy", etc. denote both straight and branched hydrocarbon groups; but reference to an individual radical such as "propyl" embraces only the straight chain radical, a branched chain isomer such as "isopropyl" is specifically referenced.

"Organometallic compound" is an organic compound comprised of a metal attached directly to carbon (R-M).

"Coordination compound" is a compound formed by the union of a central metal atom or ion with ions or molecules called ligands or complexing agents.

"Ligand" or a "complexing agent" is a molecule, ion or atom that is attached to the central metal atom or ion of a coordination compound.

"Monodentate ligand" is a ligand having a single donor atom coordinated to the central metal atom or ion.

"Bidentate ligand" is a ligand having two donor atoms coordinated to the same central metal atom or ion.

"Vanadocene" is a compound having a central vanadium metal ion coordinated with at least two cyclopentadiene groups.

It will be appreciated that compounds of the invention having a chiral center may exist in and be isolated in optically active and racemic forms. Some

compounds may exhibit polymorphism. It is to be understood that the present invention encompasses any racemic, optically-active, polymorphic, or stereoisomeric form, or mixtures thereof, of a compound of the invention, which possess the useful properties described herein. Methods to prepare optically active forms are known, for example, by resolution of the racemic form by recrystallization techniques, by synthesis from optically-active starting materials, by chiral synthesis, or by chromatographic separation using a chiral stationary phase. Methods for determining anti-mitotic and anti-meiotic activity of a compound are known, for example, using the standard tests described herein, or other known tests.

Specific values listed below for radicals, substituents, and ranges, are for illustration only; they do not exclude other defined values or other values within defined ranges for the radicals and substituents.

For example, (C₁-C₆) alkyl can be methyl, ethyl, propyl, isopropyl, butyl, iso-butyl, sec-butyl, pentyl, 3-pentyl, or hexyl; (C₁-C₃) alkyl can be methyl, ethyl or propyl; halo(C₁-C₃) alkyl can be iodomethyl, bromomethyl, chloromethyl, fluoromethyl, trifluoromethyl, 2-chloroethyl, 2-fluoroethyl, 2,2,2-trifluoroethyl, or pentafluoroethyl; (C₁-C₃) alkoxy can be methoxy, ethoxy, or propoxy; and (C₂-C₆)alkanoyloxy can be acetoxy, propanoyloxy, butanoyloxy, isobutanoyloxy, pentanoyloxy, or hexanoyloxy.

The following glossary of vanadium compounds is provided to clarify terms used throughout the specification and provides a listing of exemplary vanadium compounds useful in the method invention:

25

Group A: Vanadocene dihalides

VDC	Vanadocene dichloride (Cp ₂ VCl ₂)
VMDC	Bis (methyl cyclopentadienyl) vanadium dichloride [(MeCp) ₂ VCl ₂]
VDB	Vanadocene dibromide (Cp ₂ VBr ₂)
30 VDI	Vanadocene diiodide (Cp ₂ VI ₂)

Group B: Vanadocene di-pseudohalides

	VDA	Vanadocene diazide [$\text{Cp}_2\text{V}(\text{N}_3)_2$]
	VDCN	Vanadocene dicyanide ($\text{Cp}_2\text{V}(\text{CN})_2$)
5	VDOCN	Vanadocene dioxycyanate ($\text{Cp}_2\text{V}(\text{OCN})_2$)
	VDSCN	Vanadocene dithiocyanate ($\text{Cp}_2\text{V}(\text{SCN})_2$)
	VDSecN	Vanadocene diselenocyanate ($\text{VCp}_2(\text{SeCN})_2$)

Group C: Vanadocene disubstituted derivatives

10	VDT	Vanadocene ditriflate ($\text{Cp}_2\text{V}(\text{O}_3\text{SCF}_3)_2$)
	VDCO	Vanadocene monochloro oxycyanate ($\text{Cp}_2\text{V}(\text{OCN})(\text{Cl})$)
	VDFe	Vanadocene monoacetonitrilo monochloro tetrachloro ferrate ($\text{Cp}_2\text{VClNCCH}_3$) FeCl_4

Group D: Chelated Vanadocene Complexes

15	VDacac	Vanadocene acetylacetonato monotriflate ($\text{Cp}_2\text{V}(\text{CH}_3\text{COCH}_2\text{COCH}_3)(\text{O}_3\text{SCF}_3)$)
	VDBPY	Vanadocene bipyridino ditriflate ($\text{CP}_2\text{V}(\text{C}_{10}\text{H}_8\text{N}_2)(\text{O}_3\text{SCF}_3)_2$)
	VDHfacac	Vanadocene hexafluoro acetylacetonato monotriflate $\text{Cp}_2\text{V}(\text{CF}_3\text{COCH}_2\text{COCF}_3)(\text{O}_3\text{SCF}_3)$
20	VDH	Vanadocene acethydroxamato monotriflate ($\text{Cp}_2\text{V}(\text{CH}_3\text{CON}(\text{O})\text{H})(\text{O}_3\text{SCF}_3)$)
	VDPH	Vanadocene N-phenyl benzohydroxamato monotriflate ($\text{Cp}_2\text{V}(\text{C}_6\text{H}_5\text{CON}(\text{O})\text{C}_6\text{H}_5)(\text{O}_3\text{SCF}_3)$)

25

Group E. Oxovanadium Compounds

$[(\text{VO}(\text{phen}))]$ = (diaqua)(1,10-phenanthroline)oxovanadium (IV) sulfate;

$[(\text{VO}(\text{phen})_2)]$ = (aqua)bis(1,10-phenanthroline)oxovanadium (IV) sulfate;

[VO(Me₂-phen)] = (diaqua)(4,7-dimethyl-1,10-phenanthroline)oxovanadium (IV) sulfate;

[VO(Me₂-phen)₂] = (aqua)bis(4,7-dimethyl-1,10-phenanthroline)oxovanadium (IV) sulfate;

5 [VO(Cl-phen)] = (diaqua)(5-chloro-1,10-phenanthroline)oxovanadium (IV) sulfate;

[VO(Cl-phen)₂] = (aqua)bis(5-chloro-1,10-phenanthroline)oxovanadium (IV) sulfate;

[VO(bipy)] = (diaqua)(2,2'-bipyridyl)oxovanadium (IV) sulfate;

[VO(bipy)₂] = (aqua)bis(2,2'-bipyridyl)oxovanadium (IV) sulfate;

[VO(Me₂-bipy)] = (diaqua)(4,4'-dimethyl-2,2'-bipyridyl)oxovanadium (IV)sulfate;

10 [VO(Me₂-bipy)₂] = (aqua)bis(4,4'-dimethyl-2,2'-bipyridyl)oxovanadium (IV) sulfate;

[VO(Br,OH-acph)₂] = bis(5'-bromo-2'-hydroxyacetophenone)oxovanadium (IV).

Unless otherwise indicated, the following abbreviations are used throughout the remainder of the disclosure:

15 Cp, cyclopentadienyl

Cp⁻, cyclopentadienyl anion

acac, acetylacetonate

Bpy, 2,2' Bipyridine

Hfacac, hexafluoroacetylacetonate

20 Cat, catecholate

Dtc, diethyl dithio carbamate

Phen, penanthroline

PH, N-phenyl benzohydroxamic acids

H, acethydroxamic acid

25 OTf, trifluoromethane sulphonate

THF, tetrahydrofuran

DMSO, dimethyl sulfoxide

CH₃CN, acetonitrile

CH₂Cl₂, dichloromethane


d-d, laportte spin forbidden transitions

5 LMCT, ligand to metal charge transfer transitions


p-p*, intraligand charge transfer transitions

10 The present invention concerns vanadium compounds, and the finding that such compounds have potent and selective anti-mitotic activity, and are particularly active and stable agents for inhibiting angiogenesis. As such, these compounds are useful to treat or prevent disorders such as diabetic retinopathy, hemangiomas, cancers with abnormal blood vessel supply, restenosis following vascular injury, and the like.

15 Vanadium (IV) compounds for use in this invention are as shown in formula I and formula II:



(I)




(II)

20 where R₁ and R₂ are each independently a monodentate ligand or together form a bidentate ligand; R₃ and R₄ are each independently a monodentate ligand or together form a bidentate ligand; and R₅ is a monodentate ligand, or is absent.

25 Suitable monodentate ligands include monodentate ligands are selected from the group consisting of halo, OH₂, O₃SCF₃, N₃, CN, OCN, SCN, SeCN, and a cyclopentadienyl ring, wherein each cyclopentadienyl ring is optionally substituted with one or more (C₁-C₃)alkyl. Suitable bidentate ligands are selected from the group consisting of acac, Bpy, Hfacac, Cat, Dtc, PH, H, and Phen, or derivatives thereof. The bidentate ligands may be substituted, for example, with one or more (C₁-C₃) alkyl, halo,

(C₁-C₃) alkoxy, and halo (C₁-C₃) alkyl, and derivatives thereof. Halo is chloro, bromo, or iodo, and preferably is chloro.

In one embodiment, a useful vanadium compound has the following structure:




5

where R₁ and R₂ are each independently a monodentate ligand or together form a bidentate ligand; and R₃ and R₄ are each independently a cyclopentadienyl ring, wherein each cyclopentadienyl ring is optionally substituted with one or more (C₁-C₃)alkyl. In some preferred embodiments, R₁ and R₂ are each independently a monodentate ligand selected from the group consisting of halo, OH₂, O₃SCF₃, N₃, CN, OCN, SCN, SeCN, and a cyclopentadienyl ring, where each cyclopentadienyl ring is optionally substituted with one or more (C₁-C₃)alkyl. Preferably, R₁ and R₂ are halo, and more preferably chloro. In some other embodiments, R₁ and R₂ together form a bidentate ligand selected from the group consisting of acac, Bpy, Hfacac, Cat, Dtc, PH, H, and Phen, or derivatives thereof. Preferably, the bidentate ligand is acac, or derivatives thereof.

Some specific examples of compounds of formula I are: VCp₂Cl₂ (VDC), VCp₂Br₂, VCp₂I₂, VCp₂(N₃)₂, VCp₂(CN)₂, VCp₂(NCO)₂, VCp₂(NCO)Cl, VCp₂(NCS)₂, VCp₂(NCSe)₂, VCp₂Cl (CH₃CN)(FeCl₄), VCp₂(O₃SCF₃)₂, V(MeCp)₂Cl₂, V(Me₅Cp)₂Cl₂, VCp₂(acac) (VDacac), VCp₂(hf-acac), VCp₂(bpy), VCp₂(cat), VCp₂(dtc), VCp₂PH, or VCp₂H. Two particularly useful vandocene compounds are VDC and VDacac.

Examples of useful oxovanadium compounds of formula II include the compound having the following structure:



5

where R₁ and R₂ are each independently a monodentate ligand or together form a bidentate ligand; R₃ and R₄ together form a bidentate ligand; and R₅ is a monodentate ligand, or is absent. In some preferred embodiments, R₁ and R₂ are each independently a monodentate ligand selected from the group consisting of halo, OH₂, O₃SCF₃, N₃, CN, OCN, SCN, SeCN, and a cyclopentadienyl ring, wherein each cyclopentadienyl ring is optionally substituted with one or more (C₁-C₃)alkyl, and R₃ and R₄ together form a bidentate ligand selected from the group consisting of acac, Bpy, Hfacac, Cat, Dtc, PH, H, and Phen, or derivatives thereof. In embodiments where R₁ and R₂ together form a bidentate ligand, the bidentate ligand is preferably selected from the group consisting of acac, Bpy, Hfacac, Cat, Dtc, PH, H, and Phen, or a derivative thereof.

15

Specific compounds of formula II include [VO(phen)], [VO(phen)₂], [VO(Me₂-phen)], [VO(Me₂-phen)₂], [VO(Cl-phen)], [VO(Cl-phen)₂], [VO(bipy)], [VO(bipy)₂], [VO(Me₂-bipy)], [VO(Me₂-bipy)₂], and [VO(Br,OH-acph)₂].

20

The vandocene compounds can be used as anti-angiogenesis and anti-mitosis agents. In some embodiments, such compounds are used in the treatment of disorders in animals, and in particular in the treatment of diabetic retinopathy, hemangiomas, cancers with abnormal blood vessel supply, restenosis following vascular injury, and the like.

25

In such cases, the compounds both inhibit angiogenesis, and act as an anti-mitotic agent within the cells. In this manner the compounds of the present invention

are acting in a dual function, to both reduce the blood supply and to disrupt mitosis in dividing cells.

Administration of the compounds as salts may be appropriate. Examples
5 of acceptable salts include alkali metal (for example, sodium, potassium or lithium) or alkaline earth metal (for example calcium) salts, however, any salt that is non-toxic and effective when administered to the animal being treated is acceptable.

Acceptable salts may be obtained using standard procedures well known
in the art, for example by reacting a sufficiently acidic compound with a suitable base
10 affording a physiologically acceptable anion.

The compositions of the invention can be formulated as pharmaceutical
compositions and administered to an animal host, such as a human patient in a variety
of forms adapted to the chosen route of administration, *i.e.*, orally or parenterally, by
intravenous, intramuscular, topical or subcutaneous routes.

Thus, the present compounds may be systemically administered, *e.g.*,
15 orally, in combination with a pharmaceutically acceptable vehicle such as an inert diluent or an assimilable edible carrier. They may be enclosed in hard or soft shell gelatin capsules, may be compressed into tablets, or may be incorporated directly with the food of the patient's diet. For oral therapeutic administration, the active compound
20 may be combined with one or more excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 0.1% of active compound. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 2 to about 60% of the weight of a given unit dosage
25 form. The amount of active compound in such therapeutically useful compositions is such that an effective dosage level will be obtained. When administered orally, the compositions of the invention can preferably be administered in a gelatin capsule.

The tablets, troches, pills, capsules, and the like may also contain the
following: binders such as gum tragacanth, acacia, corn starch or gelatin; excipients
30 such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch,

alginic acid and the like; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, fructose, lactose or aspartame or a flavoring agent such as peppermint, oil of wintergreen, or cherry flavoring may be added. When the unit dosage form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier, such as a vegetable oil or a polyethylene glycol. Various other materials may be present as coatings or to otherwise modify the physical form of the solid unit dosage form. For instance, tablets, pills, or capsules may be coated with gelatin, wax, shellac or sugar and the like. A syrup or elixir may contain the active compound, sucrose or fructose as a sweetening agent, methyl and propylparabens as preservatives, a dye and flavoring such as cherry or orange flavor. Of course, any material used in preparing any unit dosage form should be pharmaceutically acceptable and substantially non-toxic in the amounts employed. In addition, the active compound may be incorporated into sustained-release preparations and devices.

The compositions of the invention may also be administered intravenously or intraperitoneally by infusion or injection. Solutions of the active composition can be prepared in water, optionally mixed with a nontoxic surfactant. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, triacetin, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

The pharmaceutical dosage forms suitable for injection or infusion can include sterile aqueous solutions or dispersions or sterile powders comprising the active ingredient which are adapted for the extemporaneous preparation of sterile injectable or infusible solutions or dispersions, optionally encapsulated in liposomes. In all cases, the ultimate dosage form should be sterile, fluid and stable under the conditions of manufacture and storage. The liquid carrier or vehicle can be a solvent or liquid dispersion medium comprising, for example, water, ethanol, a polyol (for example, glycerol, propylene glycol, liquid polyethylene glycols, and the like), vegetable oils, nontoxic glyceryl esters, and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the formation of liposomes, by the maintenance of the required particle size in the case of dispersions or by the use of surfactants. The

prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, buffers or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions are prepared by incorporating the active composition in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filter sterilization. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and the freeze drying techniques, which yield a powder of the active ingredient plus any additional desired ingredient present in the previously sterile-filtered solutions.

For topical administration, the present compositions may be applied in pure form, i.e., when they are liquids. However, it will generally be desirable to administer them to the skin as compositions or formulations, in combination with a dermatologically acceptable carrier, which may be a solid or a liquid.

Useful solid carriers include finely divided solids such as talc, clay, microcrystalline cellulose, silica, alumina and the like. Useful liquid carriers include water, alcohols or glycols or water-alcohol/glycol blends, in which the present compounds can be dissolved or dispersed at effective levels, optionally with the aid of non-toxic surfactants. Adjuvants such as fragrances and additional antimicrobial agents can be added to optimize the properties for a given use. The resultant liquid compositions can be applied from absorbent pads, used to impregnate bandages and other dressings, or sprayed onto the affected area using pump-type or aerosol sprayers.

Thickeners such as synthetic polymers, fatty acids, fatty acid salts and esters, fatty alcohols, modified celluloses or modified mineral materials can also be employed with liquid carriers to form spreadable pastes, gels, ointments, soaps, and the like, for application directly to the skin of the user.

Examples of useful dermatological compositions which can be used to deliver the compounds of formula I to the skin are known to the art; for example, see Jacquet et al. (U.S. Pat. No. 4,608,392), Geria (U.S. Pat. No. 4,992,478), Smith et al. (U.S. Pat. No. 4,559,157) and Wortzman (U.S. Pat. No. 4,820,508).

5 Useful dosages of the compounds of the present invention can be determined by comparing their *in vitro* activity, and *in vivo* activity in animal models. Methods for the extrapolation of effective dosages in mice, and other animals, to humans are known to the art; for example, see U.S. Pat. No. 4,938,949.

10 Generally, the concentration of the compositions of the invention in a liquid composition, such as a lotion, will be from about 0.1-50 wt-%, preferably from about 0.5-5 wt%. The concentration in a semi-solid or solid composition such as a gel or a powder will be about 0.1-5 wt-%, preferably about 0.5-2.5 wt-%.

15 The amount of the composition required for use in treatment will vary not only with the particular salt selected but also with the route of administration, the nature of the condition being treated, and the age and condition of the patient, and will ultimately be at the discretion of the attendant physician or clinician.

20 In general, however, a suitable dose will be in the range of from about 0.1 to about 150 mg/kg, e.g., from about 10 to about 75 mg/kg of body weight per day, such as 3 to about 50 mg per kilogram body weight of the recipient per day, preferably in the range of 1 to 100 mg/kg/day, most preferably in the range of 5 to 20 mg/kg/day.

The compositions are conveniently administered in unit dosage form; for example, containing 5 to 1000 mg, conveniently 10 to 750 mg, most conveniently, 50 to 500 mg of active ingredient per unit dosage form.

25 Ideally, the active ingredient should be administered to achieve peak plasma concentrations of the active compound of from about 0.5 to about 75 μ M, preferably, about 1 to 50 μ M, most preferably, about 2 to about 30 μ M. This may be achieved, for example, by the intravenous injection of a 0.05 to 5% solution of the active ingredient, optionally in saline, or orally administered as a bolus containing about 1-100 mg of the active ingredient. Desirable blood levels may be maintained by

continuous infusion to provide about 0.01-5.0 mg/kg/hr or by intermittent infusions containing about 0.4-15 mg/kg of the active ingredient(s).

The desired dose may conveniently be presented in a single dose or as divided doses administered at appropriate intervals, for example, as two, three, four or
5 more sub-doses per day. The sub-dose itself may be further divided, e.g., into a number of discrete loosely spaced administrations; such as multiple inhalations from an insufflator or by application of a plurality of drops into the eye.

Following i.m. administration, the compositions of the invention enter the blood stream within about 10-15 minutes and reach a maximum concentration in the
10 blood within one hour of administration, at which point they can be found throughout the circulatory related organs.

Note, for the examples that follow, Tetraphenyl borate, trifluoromethanesulfonato and acetyl acetone were purchased from Aldrich Chemical Co. (Milwaukee, WI). Other reagents used were of commercially available reagent
15 grade quality, and unless otherwise stated, all solvents were used as received from Aldrich (Sure Seal bottle, < 0.005% water). Tetrahydrofuran was dried by distillation over sodium. Infrared spectral data were recorded using a FT-Nicolet model Protege 460 and taken as a KBr pellet. Frequencies were generally in the range of 4000-500 cm^{-1} . UV-Vis spectra were recorded in a quartz cell on a Beckman Model DU 7400
20 spectrophotometer and the spectral bands were registered in the the 250 - 800 nm range. Electron paramagnetic resonance (EPR) spectra were recorded in standard PBS buffer on a Bruker ESP 300 EPR spectrophotometer (9.64 GHz) using 4102 standard cavity. The g values were calibrated with a Varian strong pitch (0.1% in KCl) standard (g value 2.0028). The samples (in PBS) for EPR spectral analysis were studied in Willmad WG-
25 814 standard TE_{102} aqueous cell cavity (0.3 mm inner path length) to minimize the dielectric loss. Hyde, Rev Sci Instrum, 43: 629, 1972. Magnetic moments were determined by Evans method in CDCl_3 on a Varian 300 MHz FTNMR spectrometer. (Evans, *J Chem Soc*, 2003, 1959).

In some embodiments, the compounds of the invention can be
30 administered before, during and/or after a vascular injury. Vascular injuries occur in

human patients, for example, after medical procedures such as coronary angioplasty. These percutaneous procedures are conducted on patients with stable angina with single vessel disease, as well as those with multivessel disease, total occlusion, complex lesions, unstable angina and acute myocardial infarction. Several procedures in
5 common use today can result in vessel injury, and would benefit from the method of the invention. These include balloon angioplasty, vessel stents, rotational and directional atherectomy, and laser angioplasty.

In a preferred embodiment, a patient is pretreated a vanadium compound such as VDacac at least one to three days before the treatment or procedure which is
10 known to induce vascular injury. For example, it is preferred that administration of the vanadium compound is performed one to three days before a medical procedure such as PTCA. Delivery of the vanadium compound preferably continues after the vascular injury or medical procedure up to a period of about 2 weeks to 6 months post injury. The compound is preferably administered to the patient for a period of about 1 month to
15 3 months post injury. It is believed that the period 1 month to 3 months is the time at which restenosis formation peaks after a vascular injury.

However, often the pretreatment option is not available to patients in an emergency situation. An emergency situation may arise requiring a medical procedure that causes vascular injury. In that situation, the vanadium compound can be
20 administered during the procedure and/or after the procedure. The compound can be administered for a period after injury of 2 weeks to about 6 months, preferably at least 1 to 3 months.

The administration of the vanadium compound ameliorates or prevents development of restenosis. Some level of neointima hyperplasia can still be present in
25 those patients treated with vanadium compounds, but the formation of neointima is significantly ameliorated compared to untreated patients. Treatment course and dose of the vanadium compounds can be adjusted in the patient if the clinical signs indicate a need for increased dose or treatment time.

Methods for analysis of restenosis are known, and include, for example,
30 quantitative coronary angiography (QCA), where narrowing of a vessel is visualized by

injection of a visualizing dye. Another method for analyzing restenosis is by intravascular ultrasound imaging (IVUS). By inserting the ultrasound probe into a vessel, the diameter of the vessel, as well as the type and extent of lesions in the vessel can be analyzed.

5

EXAMPLES

The present invention may be better understood with reference to the accompanying examples that are intended for purposes of illustration only and should not be construed to limit the scope of the invention, as defined by the claims appended hereto.

10

Example 1

Synthesis of Vanadium Compounds

Vanadium compounds useful in the invention may be prepared by known methods, as described, for example, in published PCT Applications WO99/36063; WO 00/27389; and WO 00/35930. For example, VDC (VCp_2Cl_2) and $[\text{VCp}_2(\text{acac})](\text{CF}_3\text{SO}_3)$; (VDacac) were prepared by following literature procedures (Wilkinson et al., *J. Am. Chem. Soc.*, 76: 4281-4284, 1954; Doyle et al., *Inorg. Chem.*, 7: 2479-2484, 1968) and purity was confirmed by ^1H NMR, IR spectroscopy, and elemental analysis.

20

Example 2

Inhibition of Angiogenesis

The present example illustrates that vanadium compounds such as $[\text{VCp}_2(\text{acac})](\text{CF}_3\text{SO}_3)$ are effective inhibitors of angiogenesis.

25

Chick Embryo Chorioallantoic (CAM) Assay

Inhibition of embryonic angiogenesis was determined using a bioassay system involving CAMs of growing chick embryos, as previously described (Nguyen et al, *Microvascular Research*, 47:31, 1994; Auerbach et al, *Developmental Biology*, 41:391,

30

1974). Fertilized white Leghorn chicken eggs were received at day 3 of incubation from the University of Minnesota Poultry Research Center, St. Paul, MN. The following procedures took place in a sterile laminar flow hood. The eggs were wiped down with 70% isoprpyl alcohol and allowed to air dry. The eggs were wiped with
 5 Betadine and placed in a horizontal position for approximately 5 minutes and allowed to dry. The eggs were cracked and placed into sterile 100 x 20 mm² Petri dishes (Fischer, Itasca, IL) and transferred to a 37°C humidified incubator (1.5% CO₂). Pellets containing 1-100 µg [VCp₂(acac)](CF₃SO₃) in DMSO and 0.5% methylcellulose (Sigma, St. Louis, MO) were prepared by pipetting 10 µl of the compound, using a
 10 positive displacement pipette, onto sterile TeflonTMPFA Petri dishes (VWR, Chicago, IL) and drying in a vacuum desiccator at ambient temperature. The pellets containing the indicated dose of [VCp₂(acac)](CF₃SO₃) were implanted on the outer third of a 4-6 day old CAM surface, generally between two branches of a prominent blood vessel. The eggs were returned to a humidified 1.5% CO₂/37°C incubator.

15 Inhibition of angiogenesis was assessed 48 hours after implantation by measuring the avascular zone in the CAM beneath the pellet, followed by photographic documentation of the CAM. Pellets containing 200 µg suramin (Calbiochem, La Jolla, CA) served as a positive control. The methylcellulose disk alone with no added compounds served as a negative control. Significant inhibition of angiogenesis was
 20 defined as the presence of an avascular zone of at least 3 mm in diameter around the methylcellulose disk.

Results

The effect of [VCp₂(acac)](CF₃SO₃) on angiogenesis was examined in standard
 25 chorioallantoic membrane (CAM) assays using 4-6 day old chick embryos. [VCp₂(acac)](CF₃SO₃) in methylcellulose disks at doses ranging from 1 µg to 100 µg rapidly inhibited angiogenesis and produced within one hour a large avascular zone in the CAM around the methylcellulose disk, whereas control embryos implanted with empty discs did not develop avascular zones. The average size of the avascular zone
 30 increased with increasing dose of the vanadium compound.

The data indicates that the vanadium compound [VCp₂(acac)](CF₃SO₃) is a potent inhibitor of angiogenesis.

VDacac (mg)	Avascular Zone (mm ²)
1	7.0 ± 1.5
10	38.4 ± 16.6
100	92.4 ± 24.9

5

Example 3

Inhibition of Mitosis

10 This example illustrates that the vanadium compound [VCp₂(acac)](CF₃SO₃) is an effective inhibitor of embryonic development of Zebra fish, and in particular, is a potent inhibitor of mitosis.

Zebra Fish and Embryos

15 The adult wild type ZF were maintained generally according to the “Zebrafish Book” recommendations (Westerfield, *The Zebrafish Book*, 1993, 2d Edition Univ. of Oregon Press, Eugene). Males and females were kept in 10 Gallon tanks (70 fish per tank) with a constant slow flow of conditioned water at 26 °C and a controlled 14 hour day / 10 hour night cycle. Adult fish were fed twice a day with live brine shrimp (Ocean Star International, Snowville, UT) and each group of fish was bred once in two
20 weeks. The embryos were obtained through (a) natural spawning at 28.5 °C in the breeding tanks with a netted false bottom or (b) fertilization in vitro using eggs and milt collected from the mature females and males anesthetized with Tricaine (Sigma, St.Louis, MO), as described (Westerfield, *The Zebrafish Book*, *Supra*).

25 **Zebrafish Embryo Model System**

ZF eggs were removed from their chorions by mild digestion in 1% Trypsin-EDTA (Sigma, St.Louis, MO) for 10 minutes at 28.5°C (Standard temperature - ST), washed three times in "egg water" and twice in "embryonic medium" (EM), according to recommendations (Westerfield, *The Zebrafish Book, Supra*). The dechorionated two-cell stage cleaving eggs were transferred to the 24-well plastic cell culture plates (Costar, Cambridge, MA) filled with EM. Dechorionated embryos (10-12 per well) were exposed to the drugs at a constant ST for 0.5-24 hours. The final volume of the media in each well was 500 µL. The compound was tested at concentrations ranging from 10 µM to 4 mM. First dissolved in DMSO, it was then diluted serially with the incubation medium. The final concentration of DMSO in the wells was 1.2 %. The sham-treated control embryos were incubated in EM in the presence of 1.2% DMSO.

Microinjections were performed with the help of a SMZ-10A stereo microscope (Nicon, Melville, NY) and transjector 5246 (Eppendorf Scientific Inc, Westbury, N.Y. 11590) at RT. The eggs in chorions were restrained in agar grooves filled with EM, and the drug containing solution was introduced into the cytoplasm of one of the blastomeres of the two-cell stage eggs through a micropipette with a splinted sharp tip of 2-3 µm in diameter. All microinjections were performed under visual control. Drugs for injection were dissolved in Hank's Balanced Salt Saline (Gibco, Rockville, MD) with 10% DMSO. The volume of the solution injected into each blastomere was approximately 2 nl. For every drug, 40-50 embryos were injected in 30 minutes of one series of experiment. After the treatment the embryos were transferred back to ST and studied as above.

Observations of cell division and development of the ZF embryos were carried out using a SMZ-10A stereo microscope (Nicon, Melville, NY), once every 30 minutes within the first 3 hours of incubation and at 6, 12 and 24 hours, as well. The drug effect was considered to be revealed when all embryos from one well were affected in a characteristic manner in 3 independent experiments. The stereo microscope was fitted with a specially designed transparent heating tray in order to keep embryos at ST during observations. Pictures of the embryos were taken with a H-III Photomicrographic System (Nicon, Melville, NY) using Ektachrome 64X film (Kodak, Rochester, NY).

Cytotoxicity MTT Assays

The cytotoxicity of [VCp₂(acac)](CF₃SO₃) against human cancer cell lines was performed as described previously (Ghosh et al, *Clinc Canc Res*, 6:1536, 2000) using the MTT (3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyl tetrazolium bromide) assay (Boehringer Mannheim Corp., Indianapolis, IN). Briefly, exponentially growing tumor cells were seeded into a 96-well plate at a density of 2.5×10^4 cells/well and incubated [VCp₂(acac)](CF₃SO₃) or 0.1% DMSO in PBS concentrations ranging from 0.1 to 250 μ M. Following incubation with drug for 48 hours at 37°C, to each well, 10 μ l of MTT (0.5 mg/ml final concentration) was added and the plates were incubated at 37°C for 4 hours to allow MTT to form formazan crystals by reacting with metabolically active cells. The formazan crystals were solubilized overnight at 37°C in a solution containing 10% SDS in 0.01 M HCl. The absorbance of each well was measured in a microplate reader (Labsystems) at 540 nm and a reference wavelength of 690 nm. The percent survival and the IC₅₀ values were calculated using Graphpad Prism software version 2.0 (Graphpad Software, Inc., San Diego, CA), as described previously. Ghosh et al, *Clinc Canc Res*, 6:1536, 2000.

Immunocytochemistry and Confocal Microscopic Analysis

At the appropriate time points coverslips containing BT-20 cells were fixed in -20° C methanol for 15 minutes followed by 15 minutes incubation in phosphate buffered saline + 0.1% Triton X-100 (PBS-Tx). Coverslips were next incubated with a primary antibody recognizing α -tubulin (Sigma, St.Louis MO) for 40 minutes in a humidified chamber at 37° C. Coverslips were washed for 15 minutes in PBS-Tx followed by a 40 minutes incubation with a fluorescently labeled secondary antibody (Jackson ImmunoResearch, West Grove, PA). The coverslips were again rinsed in PBS-Tx and incubated with 5 μ M Toto-3 (Molecular Probes, Eugene OR) for 20 minutes to label the DNA. Coverslips were immediately inverted onto slides in Vectashield

(Vector Labs, Burlingame, NH) to prevent photobleaching, sealed with nail polish and stored at 4° C.

Slides were examined using a Bio-Rad MRC-1024 Laser Scanning Confocal Microscope mounted on a Nikon Eclipse E800 upright microscope with high numerical aperture objectives. Slides were examined to determine the percentage of cells in interphase vs. mitosis and representative images were taken. Digital data was processed using Laserssharp (Bio-Rad, Hercules, CA) and Photoshop (Adobe Systems, Mountain View, CA) software and printed on a Pictography printer (Fuji Photo Elmsford, NY)

10 Results

Embryonic development of the ZF (*Danio rerio*) is thoroughly studied and staged (Westerfield, *The Zebrafish Book*, 1993, 2d Edition (Univ of Oregon Press, Eugene); Benyumov et al, *Rus. J. Dev. Biol.*, 26(2):132, 1995; Kimmel et al, *Dev. Dynam.*, 203:253, 1995). In a meroblastic egg, cell divisions are rapid and occur after the ooplasmic segregation on the animal pole of the egg cell, resulting within the first 3 hours of development in generation of a multicellular blastula comprised of several thousands of cells. The first series of cell divisions at the initial cleavage stage are approximately synchronous, only 15 minutes apart, and each set of the dividing blastomeres is characterized by a distinct pattern of cellular localization. This remarkable proliferation rate of undifferentiated eukaryotic vertebrate cells makes the ZF embryo an attractive experimental model for evaluation of new agents for anti-proliferative activity. In order to determine if the vanadium compounds such as [VCp₂(acac)](CF₃SO₃) could affect cell division, the effect on embryonic development of Zebra fish was examined.

As discussed above, the embryos were dechorionated with Trypsin-EDTA and incubated in embryonic medium containing either vehicle (1.2% DMSO) alone or vehicle with [VCp₂(acac)](CF₃SO₃) at different concentrations. At standard temperature of 28.5°C, two-cell stage control embryos exposed to the vehicle alone reached 4-cell, 8-cell, and 64-cell stages within 15 minutes, 30 minutes, and 75 minutes,

respectively. Within 3.5 hours post fertilization, these embryos developed into a high blastula and underwent gastrulation approximately 2.5 hours later.

[VCp₂(acac)](CF₃SO₃) inhibited cell division in a concentration-dependent fashion. At 0.6 mM, it caused cell division block at the 8-16 cell stage of embryonic development followed by total cell fusion and developmental arrest. At lower concentrations, cell proliferation continued but was abnormal. Specifically, within 120 minutes of incubation, cell division in the treated embryos was observed only on top of the blastodisc and whereas the cells on periphery and cells adjacent to the yolk were totally fused. The treated eggs did not form a compact blastoderm and developed no further.

The anti-proliferative effects of [VCp₂(acac)](CF₃SO₃) microinjected into the cytoplasm of two-cell stage ZF eggs was next examined. At the lowest dose of 1 pmol/embryo, [VCp₂(acac)](CF₃SO₃) resulted in the formation of a blastocoel-like cavity in the blastoderm of late blastulae and early gastrulae and disrupted further gastrulation. At a dose of 40 pmols/cell, VDAC slowed down cell division and resulted in deterioration of the cell localization pattern and developmental block. Thus, vanadium compounds such as [VCp₂(acac)](CF₃SO₃) are a potent anti-mitotic agents.

The anti-mitotic effects of [VCp₂(acac)](CF₃SO₃) on proliferation of human cancer cell lines was examined using MTT assays. [VCp₂(acac)](CF₃SO₃) inhibited the proliferation of the breast cancer cell lines MDA-MB-231 and BT-20 as well as the glioblastoma cell line U373 in a concentration-dependent fashion with IC₅₀ values of 9.6 μM, 25.1 μM, and 35.7 μM, respectively.

The ability of [VCp₂(acac)](CF₃SO₃) to inhibit the proliferation of human cancer cells prompted the hypothesis that this compound likely affects mitotic spindle formation. To test this hypothesis, we examined the mitotic spindles of vehicle-treated and [VCp₂(acac)](CF₃SO₃)-treated BT-20 breast cancer cells using confocal laser scanning microscopy. Whereas in vehicle-treated control cells mitotic spindles were organized as a bipolar microtubule array and the DNA was organized on a metaphase plate, [VCp₂(acac)](CF₃SO₃)-treated BT-20 cells had aberrant monopolar mitotic structures where microtubules were detected only on one side of the chromosomes and

the chromosomes were arranged in a circular pattern. These results provide a cogent explanation for the anti-miotic activity of $[\text{VCp}_2(\text{acac})](\text{CF}_3\text{SO}_3)$ against human cancer cells.


5 The data indicates that $[\text{VCp}_2(\text{acac})](\text{CF}_3\text{SO}_3)$ is a potent anti-proliferative agent, having disruptive effects on cell division during mitosis.

10 The above specification, examples and data provide a complete description of the manufacture and use of the composition of the invention. Since many embodiments of the invention can be made without departing from the spirit and scope of the invention, the invention resides in the claims hereinafter appended.


The above specification contains numerous references to literature and patent publications, each of which is hereby incorporated by reference as if fully set forth.

WE CLAIM:

1. A method for inhibiting angiogenesis comprising administering to a subject an effective angiogenesis inhibiting of a vanadium compound having the following structure:



(I)



(II)

wherein,

- R₁ and R₂ are each independently a monodentate ligand or together form a bidentate ligand;

R₃ and R₄ are each independently a monodentate ligand or together form a bidentate ligand; and

R₅ is a monodentate ligand, or is absent.

15


2. The method of claim 1, wherein each monodentate ligand is selected from the group consisting of halo, OH₂, O₃SCF₃, N₃, CN, OCN, SCN, SeCN, and a cyclopentadienyl ring, wherein the cyclopentadienyl ring is optionally substituted with one or more (C₁-C₃)alkyl, and each bidentate ligand is selected from the group consisting of acac, Bpy, Hfacac, Cat, Dtc, PH, H, Phen, or a derivative thereof.

20

3. The method of claim 2, wherein each bidentate ligand is optionally substituted with one or more of halo, (C₁-C₃) alkyl, (C₁-C₃) alkoxy, halo (C₁-C₃) alkyl, or a derivative thereof.

25

4. The method of claim 1, wherein the vanadium compound has the following structure:



wherein

R_1 and R_2 are each independently a monodentate ligand or together form a bidentate ligand; and

R_3 and R_4 are each independently a cyclopentadienyl ring, wherein each cyclopentadienyl ring is optionally substituted with one or more (C₁-C₃)alkyl.

5. The method of claim 4, wherein R_1 and R_2 are each independently a monodentate ligand selected from the group consisting of of halo, OH₂, O₃SCF₃, N₃, CN, OCN, SCN, SeCN, and a cyclopentadienyl ring, wherein each cyclopentadienyl ring is optionally substituted with one or more (C₁-C₃)alkyl.

6. The method of claim 5, wherein R_1 and R_2 are each independently halo.


7. The method of claim 6, wherein halo is chloro, bromo, or iodo.

8. The method of claim 6, wherein halo is chloro.

9. The method of claim 4, wherein R_1 and R_2 together form a bidentate ligand selected from the group consisting of acac, Bpy, Hfacac, Cat, Dtc, PH, H and derivatives thereof.

10. The method of claim 9, wherein the bidentate ligand is acac or a derivative thereof.

11. The method of claim 1, wherein the vanadium compound has the following structure:



5

wherein

R₁ and R₂ are each independently a monodentate ligand or together form a bidentate ligand;

10

R₃ and R₄ together form a bidentate ligand; and

R₅ is a monodentate ligand, or is absent.

15

12. The method of claim 11, wherein R₁ and R₂ are each independently a monodentate ligand selected from the group consisting of halo, OH₂, O₃SCF₃, N₃, CN, OCN, SCN, SeCN, and a cyclopentadienyl ring, wherein each cyclopentadienyl ring is optionally substituted with one or more (C₁-C₃)alkyl.

20

13. The method of claim 12, wherein, R₃ and R₄ together form a bidentate ligand selected from the group consisting of acac, Bpy, Hfacac, Cat, Dtc, PH, H, Phen, and derivatives thereof.

25

14. The method of claim 11, wherein R₁ and R₂ together form a bidentate ligand selected from the group consisting of acac, Bpy, Hfacac, Cat, Dtc, PH, H, Phen, and derivatives thereof.


15. The method of claim 1, wherein said vanadium compound is:

VCp₂Cl₂, VCp₂Br₂, VCp₂I₂, VCp₂(N₃)₂, VCp₂(CN)₂, VCp₂(NCO)₂,
VCp₂(NCO)Cl, VCp₂(NCS)₂, VCp₂(NCSe)₂, VCp₂Cl (CH₃CN)(FeCl₄),
VCp₂(O₃SCF₃)₂, V(MeCp)₂Cl₂, V(Me₅Cp)₂Cl₂, VCp₂(acac), VCp₂(hf-acac),
5 VCp₂(bpy), VCp₂(cat), VCp₂(dtc), VCp₂PH, or VCp₂H.


16. The method of claim 1, wherein said vanadium compound is:

[VO(phen)], [VO(phen)₂], [VO(Me₂-phen)], [VO(Me₂-phen)₂], [VO(Cl-phen)],
[VO(Cl-phen)₂], [VO(bipy)], [VO(bipy)₂], [VO(Me₂-bipy)], [VO(Me₂-bipy)₂],
10 and [VO(Br,OH-acph)₂].

17. A method for treating diabetic retinopathy in a subject, comprising administering to the subject an effective mitosis inhibiting amount of a vanadium compound having the following structure:



(I)



(II)


wherein,

R₁ and R₂ are each independently a monodentate ligand or together form a bidentate ligand;


R₃ and R₄ are each independently a monodentate ligand or together form
20 a bidentate ligand; and

R₅ is a monodentate ligand, or is absent.

18. A method for treating restenosis following coronary angioplasty in a subject, comprising administering to the subject an effective amount of a vanadium compound having the following structure:



(I)



(II)


wherein,

5 R_1 and R_2 are each independently a monodentate ligand or together form a bidentate ligand;


R_3 and R_4 are each independently a monodentate ligand or together form a bidentate ligand; and

R_5 is a monodentate ligand, or is absent.

- 10 19. A method for preventing or treating diabetic retinopathy in a subject, comprising:
administering to the subject an effective amount of administering to the subject an
effective mitosis inhibiting amount of a vanadium compound having the following
structure:



(I)



(II)

15 wherein,

R_1 and R_2 are each independently a monodentate ligand or together form a bidentate ligand;

20 R_3 and R_4 are each independently a monodentate ligand or together form a bidentate ligand; and

R_5 is a monodentate ligand, or is absent.

- 25 20. The method of claim 19, wherein the vascular injury is associated with an angioplasty procedure.

- 5

Abstract

	(a)	(b)	(c)	(d)	(e)	(f)	(g)	(h)	(i)	(j)	(k)	(l)	(m)	(n)	(o)	(p)	(q)	(r)	(s)	(t)	(u)	(v)	(w)	(x)	(y)	(z)	(aa)	(ab)	(ac)	(ad)	(ae)	(af)	(ag)	(ah)	(ai)	(aj)	(ak)	(al)	(am)	(an)	(ao)	(ap)	(aq)	(ar)	(as)	(at)	(au)	(av)	(aw)	(ax)	(ay)	(az)	(ba)	(bb)	(bc)	(bd)	(be)	(bf)	(bg)	(bh)	(bi)	(bj)	(bk)	(bl)	(bm)	(bn)	(bo)	(bp)	(bq)	(br)	(bs)	(bt)	(bu)	(bv)	(bw)	(bx)	(by)	(bz)	(ca)	(cb)	(cc)	(cd)	(ce)	(cf)	(cg)	(ch)	(ci)	(cj)	(ck)	(cl)	(cm)	(cn)	(co)	(cp)	(cq)	(cr)	(cs)	(ct)	(cu)	(cv)	(cw)	(cx)	(cy)	(cz)	(da)	(db)	(dc)	(dd)	(de)	(df)	(dg)	(dh)	(di)	(dj)	(dk)	(dl)	(dm)	(dn)	(do)	(dp)	(dq)	(dr)	(ds)	(dt)	(du)	(dv)	(dw)	(dx)	(dy)	(dz)	(ea)	(eb)	(ec)	(ed)	(ee)	(ef)	(eg)	(eh)	(ei)	(ej)	(ek)	(el)	(em)	(en)	(eo)	(ep)	(eq)	(er)	(es)	(et)	(eu)	(ev)	(ew)	(ex)	(ey)	(ez)	(fa)	(fb)	(fc)	(fd)	(fe)	(ff)	(fg)	(fh)	(fi)	(fj)	(fk)	(fl)	(fm)	(fn)	(fo)	(fp)	(fq)	(fr)	(fs)	(ft)	(fu)	(fv)	(fw)	(fx)	(fy)	(fz)	(ga)	(gb)	(gc)	(gd)	(ge)	(gf)	(gg)	(gh)	(gi)	(gj)	(gk)	(gl)	(gm)	(gn)	(go)	(gp)	(gq)	(gr)	(gs)	(gt)	(gu)	(gv)	(gw)	(gx)	(gy)	(gz)	(ha)	(hb)	(hc)	(hd)	(he)	(hf)	(hg)	(hh)	(hi)	(hj)	(hk)	(hl)	(hm)	(hn)	(ho)	(hp)	(hq)	(hr)	(hs)	(ht)	(hu)	(hv)	(hw)	(hx)	(hy)	(hz)	(ia)	(ib)	(ic)	(id)	(ie)	(if)	(ig)	(ih)	(ii)	(ij)	(ik)	(il)	(im)	(in)	(io)	(ip)	(iq)	(ir)	(is)	(it)	(iu)	(iv)	(iw)	(ix)	(iy)	(iz)	(ja)	(jb)	(jc)	(jd)	(je)	(jf)	(jg)	(jh)	(ji)	(jj)	(jk)	(jl)	(jm)	(jn)	(jo)	(jp)	(jq)	(jr)	(js)	(jt)	(ju)	(jv)	(jw)	(jx)	(jy)	(jz)	(ka)	(kb)	(kc)	(kd)	(ke)	(kf)	(kg)	(kh)	(ki)	(kj)	(kk)	(kl)	(km)	(kn)	(ko)	(kp)	(kq)	(kr)	(ks)	(kt)	(ku)	(kv)	(kw)	(kx)	(ky)	(kz)	(la)	(lb)	(lc)	(ld)	(le)	(lf)	(lg)	(lh)	(li)	(lj)	(lk)	(ll)	(lm)	(ln)	(lo)	(lp)	(lq)	(lr)	(ls)	(lt)	(lu)	(lv)	(lw)	(lx)	(ly)	(lz)	(ma)	(mb)	(mc)	(md)	(me)	(mf)	(mg)	(mh)	(mi)	(mj)	(mk)	(ml)	(mm)	(mn)	(mo)	(mp)	(mq)	(mr)	(ms)	(mt)	(mu)	(mv)	(mw)	(mx)	(my)	(mz)	(na)	(nb)	(nc)	(nd)	(ne)	(nf)	(ng)	(nh)	(ni)	(nj)	(nk)	(nl)	(nm)	(nn)	(no)	(np)	(nq)	(nr)	(ns)	(nt)	(nu)	(nv)	(nw)	(nx)	(ny)	(nz)	(oa)	(ob)	(oc)	(od)	(oe)	(of)	(og)	(oh)	(oi)	(oj)	(ok)	(ol)	(om)	(on)	(oo)	(op)	(oq)	(or)	(os)	(ot)	(ou)	(ov)	(ow)	(ox)	(oy)	(oz)	(pa)	(pb)	(pc)	(pd)	(pe)	(pf)	(pg)	(ph)	(pi)	(pj)	(pk)	(pl)	(pm)	(pn)	(po)	(pp)	(pq)	(pr)	(ps)	(pt)	(pu)	(pv)	(pw)	(px)	(py)	(pz)	(qa)	(qb)	(qc)	(qd)	(qe)	(qf)	(qg)	(qh)	(qi)	(qj)	(qk)	(ql)	(qm)	(qn)	(qo)	(qp)	(qq)	(qr)	(qs)	(qt)	(qu)	(qv)	(qw)	(qx)	(qy)	(qz)	(ra)	(rb)	(rc)	(rd)	(re)	(rf)	(rg)	(rh)	(ri)	(rj)	(rk)	(rl)	(rm)	(rn)	(ro)	(rp)	(rq)	(rr)	(rs)	(rt)	(ru)	(rv)	(rw)	(rx)	(ry)	(rz)	(sa)	(sb)	(sc)	(sd)	(se)	(sf)	(sg)	(sh)	(si)	(sj)	(sk)	(sl)	(sm)	(sn)	(so)	(sp)	(sq)	(sr)	(ss)	(st)	(su)	(sv)	(sw)	(sx)	(sy)	(sz)	(ta)	(tb)	(tc)	(td)	(te)	(tf)	(tg)	(th)	(ti)	(tj)	(tk)	(tl)	(tm)	(tn)	(to)	(tp)	(tq)	(tr)	(ts)	(tt)	(tu)	(tv)	(tw)	(tx)	(ty)	(tz)	(ua)	(ub)	(uc)	(ud)	(ue)	(uf)	(ug)	(uh)	(ui)	(uj)	(uk)	(ul)	(um)	(un)	(uo)	(up)	(uq)	(ur)	(us)	(ut)	(uu)	(uv)	(uw)	(ux)	(uy)	(uz)	(va)	(vb)	(vc)	(vd)	(ve)	(vf)	(vg)	(vh)	(vi)	(vj)	(vk)	(vl)	(vm)	(vn)	(vo)	(vp)	(vq)	(vr)	(vs)	(vt)	(vu)	(vv)	(vw)	(vx)	(vy)	(vz)	(wa)	(wb)	(wc)	(wd)	(we)	(wf)	(wg)	(wh)	(wi)	(wj)	(wk)	(wl)	(wm)	(wn)	(wo)	(wp)	(wq)	(wr)	(ws)	(wt)	(wu)	(wv)	(ww)	(wx)	(wy)	(wz)	(xa)	(xb)	(xc)	(xd)	(xe)	(xf)	(xg)	(xh)	(xi)	(xj)	(xk)	(xl)	(xm)	(xn)	(xo)	(xp)	(xq)	(xr)	(xs)	(xt)	(xu)	(xv)	(xw)	(xx)	(xy)	(xz)	(ya)	(yb)	(yc)	(yd)	(ye)	(yf)	(yg)	(yh)	(yi)	(yj)	(yk)	(yl)	(ym)	(yn)	(yo)	(yp)	(yq)	(yr)	(ys)	(yt)	(yu)	(yv)	(yw)
--	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

MERCHANT & GOULD P.C.

United States Patent Application

COMBINED DECLARATION AND POWER OF ATTORNEY

As a below named inventor I hereby declare that: my residence, post office address and citizenship are as stated below next to my name; that

I verily believe I am the original, first and sole inventor (if only one name is listed below) or a joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: VANADIUM COMPOUNDS AS ANTI-ANGIOGENIC AGENTS

The specification of which

- a. ☒ is attached hereto
b. ☐ was filed on as application serial no. and was amended on (if applicable) (in the case of a PCT-filed application) described and claimed in international no. filed and as amended on (if any), which I have reviewed and for which I solicit a United States patent.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the patentability of this application in accordance with Title 37, Code of Federal Regulations, § 1.56 (attached hereto).

I hereby claim foreign priority benefits under Title 35, United States Code, § 119/365 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on the basis of which priority is claimed:

- a. ☒ no such applications have been filed.
b. ☐ such applications have been filed as follows:

FOREIGN APPLICATION(S), IF ANY, CLAIMING PRIORITY UNDER 35 USC § 119			
COUNTRY	APPLICATION NUMBER	DATE OF FILING (day, month, year)	DATE OF ISSUE (day, month, year)
ALL FOREIGN APPLICATION(S), IF ANY, FILED BEFORE THE PRIORITY APPLICATION(S)			
COUNTRY	APPLICATION NUMBER	DATE OF FILING (day, month, year)	DATE OF ISSUE (day, month, year)

I hereby claim the benefit under Title 35, United States Code, § 120/365 of any United States and PCT international application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, § 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, § 1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. APPLICATION NUMBER	DATE OF FILING (day, month, year)	STATUS (patented, pending, abandoned)

I hereby claim the benefit under Title 35, United States Code § 119(e) of any United States provisional application(s) listed below:

U.S. PROVISIONAL APPLICATION NUMBER	DATE OF FILING (Day, Month, Year)

I hereby appoint the following attorney(s) and/or patent agent(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected herewith:

Albrecht, John W.	Reg. No. 40,481	Leon, Andrew J.	Reg. No. 46,869
Ali, M. Jeffer	Reg. No. 46,359	Leonard, Christopher J.	Reg. No. 41,940
Anderson, Gregg I.	Reg. No. 28,828	Liepa, Mara E.	Reg. No. 40,066
Batzli, Brian H.	Reg. No. 32,960	Lindquist, Timothy A.	Reg. No. 40,701
Beard, John L.	Reg. No. 27,612	Lycke, Lawrence E.	Reg. No. 38,540
Berns, John M.	Reg. No. 43,496	Mayfield, Denise L.	Reg. No. 33,732
Black, Bruce E.	Reg. No. 41,622	McDonald, Daniel W.	Reg. No. 32,044
Branch, John W.	Reg. No. 41,633	McIntyre, Jr., William F.	Reg. No. 44,921
Bremer, Dennis C.	Reg. No. 40,528	Mitchem, M. Todd	Reg. No. 40,731
Bruess, Steven C.	Reg. No. 34,130	Mueller, Douglas P.	Reg. No. 30,300
Byrne, Linda M.	Reg. No. 32,404	Nichols, A. Shane	Reg. No. 43,836
Campbell, Keith	Reg. No. P-46,597	Pauly, Daniel M.	Reg. No. 40,123
Carlson, Alan G.	Reg. No. 25,959	Phillips, Bryan K.	Reg. No. P-46,990
Caspers, Philip P.	Reg. No. 33,227	Phillips, John B.	Reg. No. 37,206
Chiapetta, James R.	Reg. No. 39,634	Prendergast, Paul	Reg. No. 46,068
Clifford, John A.	Reg. No. 30,247	Pytel, Melissa J.	Reg. No. 41,512
Coldren, Richard J.	Reg. No. 44,084	Qualey, Terry	Reg. No. 25,148
Daignault, Ronald A.	Reg. No. 25,968	Reich, John C.	Reg. No. 37,703
Daley, Dennis R.	Reg. No. 34,994	Reiland, Earl D.	Reg. No. 25,767
Dalglisch, Leslie E.	Reg. No. 40,579	Samuels, Lisa A.	Reg. No. 43,080
Daulton, Julie R.	Reg. No. 36,414	Schmaltz, David G.	Reg. No. 39,828
DeVries Smith, Katherine M.	Reg. No. 42,157	Schuman, Mark D.	Reg. No. 31,197
DiPietro, Mark J.	Reg. No. 28,707	Schumann, Michael D.	Reg. No. 30,422
Edell, Robert T.	Reg. No. 20,187	Scull, Timothy B.	Reg. No. 42,137
Epp Ryan, Sandra	Reg. No. 39,667	Sebald, Gregory A.	Reg. No. 33,280
Glance, Robert J.	Reg. No. 40,620	Skoog, Mark T.	Reg. No. 40,178
Goggin, Matthew J.	Reg. No. 44,125	Spellman, Steven J.	Reg. No. 45,124
Golla, Charles E.	Reg. No. 26,896	Stoll-DeBell, Kirstin L.	Reg. No. 43,164
Gorman, Alan G.	Reg. No. 38,472	Sumner, John P.	Reg. No. 29,114
Gould, John D.	Reg. No. 18,223	Swenson, Erik G.	Reg. No. 45,147
Gregson, Richard	Reg. No. 41,804	Tellekson, David K.	Reg. No. 32,314
Gresens, John J.	Reg. No. 33,112	Trembath, Jon R.	Reg. No. 38,344
Hamer, Samuel A.	Reg. No. 46,754	Tuchman, Ido	Reg. No. 45,924
Hamre, Curtis B.	Reg. No. 29,165	Tunheim, Marcia A.	Reg. No. 42,189
Harrison, Kevin C.	Reg. No. P-46,759	Underhill, Albert L.	Reg. No. 27,403
Hertzberg, Brett A.	Reg. No. 42,660	Vandenburgh, J. Derek	Reg. No. 32,179
Hillson, Randall A.	Reg. No. 31,838	Wahl, John R.	Reg. No. 33,044
Holzer, Jr., Richard J.	Reg. No. 42,668	Weaver, Karrie G.	Reg. No. 43,245
Johnston, Scott W.	Reg. No. 39,721	Welter, Paul A.	Reg. No. 20,890
Kadievitch, Natalie D.	Reg. No. 34,196	Whipps, Brian	Reg. No. 43,261
Karjeker, Shaukat	Reg. No. 34,049	Whitaker, John E.	Reg. No. 42,222
Kastelic, Joseph M.	Reg. No. 37,160	Wickhem, J. Scot	Reg. No. 41,376
Kettelberger, Denise	Reg. No. 33,924	Williams, Douglas J.	Reg. No. 27,054
Keys, Jeramie J.	Reg. No. 42,724	Withers, James D.	Reg. No. 40,376
Knearl, Homer L.	Reg. No. 21,197	Witt, Jonelle	Reg. No. 41,980
Kowalchyk, Alan W.	Reg. No. 31,535	Wu, Tong	Reg. No. 43,361
Kowalchyk, Katherine M.	Reg. No. 36,848	Xu, Min S.	Reg. No. 39,536
Lacy, Paul E.	Reg. No. 38,946	Zeuli, Anthony R.	Reg. No. 45,255
Larson, James A.	Reg. No. 40,443		

I hereby authorize them to act and rely on instructions from and communicate directly with the person/assignee/attorney/firm/ organization who/which first sends/sent this case to them and by whom/which I hereby declare that I have consented after full disclosure to be represented unless/until I instruct Merchant & Gould P.C. to the contrary.

Please direct all correspondence in this case to Merchant & Gould P.C. at the address indicated below:

Merchant & Gould P.C.
P.O. Box 2903
Minneapolis, MN 55402-0903



I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

2	Full Name Of Inventor	Family Name Uckun	First Given Name Fatih	Second Given Name M
0	Residence & Citizenship	City White Bear Lake	State or Foreign Country Minnesota	Country of Citizenship USA
1	Post Office Address	Post Office Address 12590 Ethan Avenue North	City White Bear Lake	State & Zip Code/Country MN 55110/USA
Signature of Inventor 201:				Date:

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398	2399	2400	2401	2402	2403	2404	2405	2406	2407	2408	2409	2410	2411	2412	2413	2414	2415	2416	2417	2418	2419	2420	2421	2422	2423	2424	2425	2426	2427	2428	2429	2430	2431	2432	2
--	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	---

§ 1.56 Duty to disclose information material to patentability.

(a) A patent by its very nature is affected with a public interest. The public interest is best served, and the most effective patent examination occurs when, at the time an application is being examined, the Office is aware of and evaluates the teachings of all information material to patentability. Each individual associated with the filing and prosecution of a patent application has a duty of candor and good faith in dealing with the Office, which includes a duty to disclose to the Office all information known to that individual to be material to patentability as defined in this section. The duty to disclose information exists with respect to each pending claim until the claim is canceled or withdrawn from consideration, or the application becomes abandoned. Information material to the patentability of a claim that is canceled or withdrawn from consideration need not be submitted if the information is not material to the patentability of any claim remaining under consideration in the application. There is no duty to submit information which is not material to the patentability of any existing claim. The duty to disclose all information known to be material to patentability is deemed to be satisfied if all information known to be material to patentability of any claim issued in a patent was cited by the Office or submitted to the Office in the manner prescribed by §§ 1.97(b)-(d) and 1.98. However, no patent will be granted on an application in connection with which fraud on the Office was practiced or attempted or the duty of disclosure was violated through bad faith or intentional misconduct. The Office encourages applicants to carefully examine:

(1) prior art cited in search reports of a foreign patent office in a counterpart application, and

(2) the closest information over which individuals associated with the filing or prosecution of a patent application believe any pending claim patentably defines, to make sure that any material information contained therein is disclosed to the Office.

(b) Under this section, information is material to patentability when it is not cumulative to information already of record or being made of record in the application, and

(1) It establishes, by itself or in combination with other information, a prima facie case of unpatentability of a claim;

(2) It refutes, or is inconsistent with, a position the applicant takes in:

(i) Opposing an argument of unpatentability relied on by the Office, or

(ii) Asserting an argument of patentability.

A prima facie case of unpatentability is established when the information compels a conclusion that a claim is unpatentable under the preponderance of evidence, burden-of-proof standard, giving each term in the claim its broadest reasonable construction consistent with the specification, and before any consideration is given to evidence which may be submitted in an attempt to establish a contrary conclusion of patentability.

(c) Individuals associated with the filing or prosecution of a patent application within the meaning of this section are:

(1) Each inventor named in the application:

(2) Each attorney or agent who prepares or prosecutes the application; and

(3) Every other person who is substantively involved in the preparation or prosecution of the application and who is associated with the inventor, with the assignee or with anyone to whom there is an obligation to assign the application.

(d) Individuals other than the attorney, agent or inventor may comply with this section by disclosing information to the attorney, agent, or inventor.

NONPROFIT ORGANIZATION
VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS
(37 C.F.R. 1.9(e) AND 1.27(d)) - NONPROFIT ORGANIZATION

I hereby declare that I am an official empowered to act on behalf of the nonprofit organization identified below:

NAME OF ORGANIZATION: Parker Hughes Institute
ADDRESS OF ORGANIZATION: 2665 Long Lake Road
Suite 330
Roseville, MN 55113

TYPE OF ORGANIZATION:

- CHECK ONE a) ☐ UNIVERSITY OR OTHER INSTITUTION OF HIGHER EDUCATION
b) ☒ TAX EXEMPT UNDER INTERNAL REVENUE SERVICE CODE (26 U.S.C. 501(a) AND 501(c)(3))
c) ☐ NONPROFIT SCIENTIFIC OR EDUCATIONAL UNDER STATUTE OF STATE OF THE UNITED STATES OF AMERICA -- (NAME OF STATE __)
(CITATION OF STATUTE __)
d) ☐ WOULD QUALIFY AS TAX EXEMPT UNDER INTERNAL REVENUE SERVICE CODE (26 U.S.C. 501(a) AND 501(c)(3)) IF LOCATED IN THE UNITED STATES OF AMERICA
e) ☐ WOULD QUALIFY AS NONPROFIT SCIENTIFIC OR EDUCATIONAL UNDER STATUTE OF STATE OF THE UNITED STATES OF AMERICA IF LOCATED IN THE UNITED STATES OF AMERICA -- (NAME OF STATE __)
(CITATION OF STATUTE __)

I hereby declare that the nonprofit organization identified above qualifies as a nonprofit organization as defined in 37 C.F.R. 1.9(e) for purposes of paying reduced fees under Section 41(a) and (b) of Title 35, United States Code, in regard to the invention, entitled VANADIUM COMPOUNDS AS ANTI-ANGIOGENIC AGENTS by inventor(s) Fatih M. Uckun described in

- a) ☒ the specification filed herewith.
b) ☐ provisional application serial no. __, filed __.
c) ☐ non-provisional application serial no. __, filed __.
d) ☐ patent no. __, issued __.

I hereby declare that rights under contract or law have been conveyed to and remain with the nonprofit organization with regard to the above identified invention.

If the rights held by the nonprofit organization are not exclusive, each individual, concern or organization having rights to the invention is listed below* and no rights to the invention are held by any person, other than the inventor, who could not qualify as an independent inventor under 37 C.F.R. 1.9(c) or by any concern which would not qualify as a small business concern under 37 C.F.R. 1.9(d) or a nonprofit organization under 37 C.F.R. 1.9(e). *NOTE: Separate verified statements are required from each named person, concern or organization having rights to the invention averring to their status as small entities. (37 C.F.R. 1.27)INSERT CO-OWNERS, IF ANY

NAME _____
ADDRESS _____
a) ☐ INDIVIDUAL b) ☐ SMALL BUSINESS CONCERN c) ☐ NONPROFIT ORGANIZATION

NAME _____
ADDRESS _____
a) ☐ INDIVIDUAL b) ☐ SMALL BUSINESS CONCERN c) ☐ NONPROFIT ORGANIZATION

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 C.F.R. 1.28(b))

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that

willful false statements and the like so made are punishable by fine or imprisonment, or both under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereof, or any patent to which this verified statement is directed.

NAME: Fatih M. Uckun
TITLE:

ADDRESS: 2665 Long Lake Road, Suite 330
Roseville, MN 55113

SIGNATURE: _____

DATE: _____

009770820

